

RHIC Perfect Fluid sQGP- a Landau Chromo-Diamagnetic Liquid

Somenath Chakrabarty

*Department of Physics, Visva-Bharati, Santiniketan 731 235,
West Bengal, India, E-mail:somenath@vbphysics.net.in*

We speculate that the quark matter produced at RHIC, behaves like a perfect fluid may be considered as Landau chromo-diamagnetic material in presence of strong color magnetic field. We have shown that in such a system the viscosity coefficient can be small enough and in the extreme case of ultra-strong color magnetic field strength, it behaves like a super fluid.

PACS numbers: 97.60.Jd, 97.60.-s, 75.25.+z

The possibility of deconfinement transition in ultra-relativistic heavy-ion collisions was predicted long ago. A lot of theoretical work have been done on the order of phase transition to quark matter and also the possible signatures of QGP formation in ultra-relativistic heavy-ion collisions. A number of models for space-time evolution of QGP phase were also proposed [1]. In the cosmological scenario, it is also expected that the universe was filled with hot and dense soup of QGP at about micro-second after big-bang [2].

In the past high energy heavy-ion collision experiments, starting from SPS energies (which is of course not too high), the theoretically predicted experimental signatures of QGP were either not observed or were found anomalous in the sense that the same signatures one could get if it were instead of QGP a hot and dense hadronic matter. The main reason behind all these ambiguities is the scale of incident energy of the heavy-ion beam.

Only recently RHIC offers the experimentalists and also the theoreticians who once predicted a lot of experimental signatures of QGP to study such a hot and dense system in the extreme situation, which was also expected to be present in the very early universe. At RHIC, when two gold nuclei moving with almost the speed of light, collide, the amount of energy that dump in the microscopic fireball is about $2.0 \times 10^4 \text{ GeV}$. The pressure generated at the time of impact of two gold nuclei is truly immense, it is about 10^{30} atmospheric pressure and the temperature inside the fireball is about trillions of degrees. Under such extreme condition, it is expected that all the (constituent) quarks of the colliding nuclei should come out of their parent hadrons and form a gas of hot and dense quark matter. Not only that, because of such high temperature and densities, an enormous number of quark anti-quark pairs are also produced from vacuum and all of them move in a self-consistently generated color field. Because of pair annihilation processes within the hot QGP system, there will be a lot of thermal gluons. This whole thing is expected to behave like a plasma of quarks (anti-quarks) and gluons, called quark-gluon plasma (QGP). This physical scenario was predicted long ago by the theoretical high-energy heavy-ion physicists or high-energy nuclear physicists.

Surprisingly, the physical picture emerging from the detectors BRAHMS, PHENIX, PHOBOS and STAR of four different experimental groups is consistent. The recorded data of sub-atomic debris coming out of the violent collisions at the four colliding points of RHIC, observed by the four detectors as mentioned above, show a complete deconfinement of quarks (constituents and sea quarks) and gluons. Most surprisingly, this hot and dense quark matter, instead of a gas, behaves like a liquid in thermal equilibrium. It exhibits a collective flow of almost a perfect liquid, i.e., a fluid of extremely low shear viscosity [3]. The elliptic flow of QGP is a direct evidence of its liquid like behaviour. Further, the elliptic pattern of flow indicates the presence of substantial pressure gradient in QGP and that the quarks and gluons which subsequently produce hadrons having anisotropic spatial distribution, show the collective nature. People are calling this quark matter system by strongly correlated QGP or in short, sQGP.

This situation conflicts with the theoretical prediction that the quark matter if produced in ultra-relativistic heavy-ion collisions will be almost ideal, weakly interacting gas. Since the hadrons from RHIC sQGP are not emitted uniformly in all possible directions, it is not acceptable that the hot quark matter is in a gas phase. On the other hand, it is seen that the produced quark matter behaves like a most perfect liquid ever observed.

There are a number of theoretical attempts and also some suggestions to explain this surprising physical scenario produced at RHIC. One of the approach is the lattice QCD calculation using high speed sophisticated computers. However, the most up to date version of lattice QCD calculation is unable to handle the rapidly changing scenario of RHIC sQGP and reproduce the jet quenching and low viscosity phenomenon [4]. It is also suggested to get some help from string theorists, people working on super-symmetry, super-gravity, quantum black holes, etc. In a recent work an anomalous viscosity [5], which is low enough compared to kinetic viscosity is introduced for expanding RHIC sQGP. It is shown using the turbulent motion of QGP in presence of self-consistently generated color magnetic fields that there can exist an anomalous viscosity of very low magnitude compared to the viscosity arising from particle collisions.

In the present article we shall try to explain the low viscosity nature of RHIC sQGP from a completely different line of thought. We assume the motion of thermal gluons and quarks (anti-quarks) in a self consistently generated

color magnetic field. Now it is well known from QED that in a charge particle system in presence of an external magnetic field, if the strength of the field exceeds some quantum critical value, which is a function of mass and magnitude of charge carried by the particle concerned, the Landau levels of the charge particles will be populated [6]. The phenomenon is called Landau diamagnetism. It is the quantum mechanical effect of strong magnetic field. The phenomenon is also known as Landau quantization (the transverse component of particle momentum gets quantized) in condensed matter physics. We expect that the same phenomenon can be possible in presence of self-consistently generated color magnetic field inside RHIC sQGP. The quarks (anti-quarks) and thermal gluons, carrying color charges will interact with the strong color magnetic field quantum mechanically and if the field strength exceeds the corresponding quantum critical limit, the Landau levels of these particles will be populated and as a consequence the RHIC sQGP will behave like a Landau chromo-diamagnetic system. To develop a formalism to obtain the shear viscosity coefficient of such a system, the first step we have followed is to choose the gauge $A_a^\mu \equiv (0, 0, xB_a, 0)$, where A_a^μ is the color electric field vector and B_a is the color magnetic field along z -axis, $a = 1, 2, \dots, 8$, the color index. Then the single particle energy for quarks (anti-quarks) is given by

$$E_{n,\alpha}(k_z) = \left[k_z^2 + m^2 + 2eB\left(n + \frac{1}{2} + \alpha\right) \right]^{1/2} \quad (1)$$

where k_z is the z -component of quark momentum, varies continuously from $-\infty$ to $+\infty$, m is the quark mass, $n = 0, 1, 2, \dots$ is called Landau quantum number, $\alpha = \pm 1/2$ the spin projection and $eB = \sum gq_a B_a$, with q_a is the color charge related to the generators of irreducible $SU(3)$ representation and g is the corresponding strength and the sum is over the color index. The quantity $[(2n+1+2\alpha)eB]^{1/2} = p_\perp$ is the orthogonal component of quark (anti-quark) momentum. Since the transverse part of momentum $p_\perp$ is changing in a discrete manner, the phenomenon is called the Landau quantization. We can similarly write down the energy eigen value for gluons, given by

$$E_{n,\alpha}(k_z) = \left[k_z^2 + 2eB\left(n + \frac{1}{2} + \alpha\right) \right]^{1/2} \quad (2)$$

where, in this case $\alpha = \pm 1$, the corresponding spin projections for two physical degrees of freedom of gluons. In the present scenario the geometrical structure of momentum space for the colored particles will also get modified. It will become cylindrical instead of usual spherical structure, with the axis along the direction of color magnetic field.

Now it is already known from QED calculations that the upper limit of Landau quantum number ($\nu_{\max}$) for quarks decreases with the increase of magnetic field strength. Therefore only low lying Landau levels will be populated for quarks (of course there will be no change for the gluonic case, which are vector bosons carrying color charges) if the strength of color magnetic field is strong enough. In the extreme situation, when $\nu_{\max}$ becomes zero, all the quarks (anti-quarks) will occupy only the zeroth Landau level. In this case the effect of color magnetic field on (color) charged particles will be most prominent.

To obtain an expression for shear viscosity coefficient for expanding RHIC sQGP, we study the transport theory of this Landau chromo-diamagnetic matter following the relativistic version of Boltzmann kinetic equation. The relevant part of which is given by [7]

$$p^\mu \partial_\mu f(x, p) = C[f] \quad (3)$$

where $f(x, p)$ is the distribution function and $C[f]$ is the collision terms, which contains the rates of all possible elementary processes. With the relaxation time approximation, we have

$$C[f] = -\frac{p^0}{\tau} [f(x, p) - f_0(p)] \quad (4)$$

where τ is the relaxation time and $f_0(p)$ is the equilibrium distribution function (Fermi distribution for quarks and anti-quarks and Bose distribution for gluons). We further make linear approximation for the non-equilibrium distribution function, which gives

$$f(x, p) = f_0(p)[1 + \chi(x, p)] \quad (5)$$

Then following de Groot [7], we have the expression for shear viscosity coefficient for quarks (anti-quarks)

$$\eta_{q\bar{q}} = \frac{g_q}{T} \sum_{\nu=0}^{\infty} (2 - \delta_{\nu 0}) \frac{eB}{4\pi^2} \int_{-\infty}^{+\infty} \frac{\tau_q}{E_\nu} dp_z (p_r p_z)^2 [f_0(p)(1 - f_0(p)) + \bar{f}_0(p)(1 - \bar{f}_0(p))] \quad (6)$$

where $g_g = (2N_f + 1)(2N_c + 1)$ is the degeneracy factor, T is the temperature of the system, $p_r = p_\perp = (2\nu eB)^{1/2}$ and $2\nu = (2n + 1 + 2s_z)$, with $\nu = 0, 1, 2, \dots$ etc.

Similarly for gluons, we have

$$\eta_g = \frac{g_g}{T} \sum_{\nu=0}^{\infty} (2 - \delta_{\nu 0}) \frac{eB}{4\pi^2} \int_{-\infty}^{+\infty} \frac{\tau_g}{E_\nu} dp_z (p_r p_z)^2 f_0(p) (1 + f_0(p)) \quad (7)$$

where the degeneracy factor $g_g = (2N_c + 1)$. It is easy to show that the zeroth Landau levels for quarks, anti-quarks and gluons are singly degenerate, whereas all other states are doubly degenerate. To obtain these expressions, we have assumed that the flow is along z -direction. However, since the expressions are symmetric with respect to p_r and p_z , there will be no qualitative or quantitative change if we assume flow along some direction on the $x - y$ plane. Now following the arguments that the relaxation rates are additive, we have

$$\frac{1}{\eta_s} = \frac{1}{\eta_{q\bar{q}}} + \frac{1}{\eta_g} \quad (8)$$

where η_s is the effective viscosity coefficient of the system. It is quite obvious from the expression for shear viscosity coefficient for quarks (anti-quarks) that in the extreme case when only the zeroth Landau level is occupied, the system behaves like a super fluid. In fig.(1), just for illustration we have shown the variation of effective viscosity coefficient with the strength of color magnetic field, normalized with respect to QED quantum critical value for electron ($4.4 \times 10^{13} \text{G}$). We have chosen two different temperatures and densities of the matter. To obtain the relaxation time, we have considered the following elementary processes: $qq \rightarrow qq$, $q\bar{q} \rightarrow q\bar{q}$, $\bar{q}\bar{q} \rightarrow \bar{q}\bar{q}$, $qg \rightarrow qg$, $\bar{q}g \rightarrow \bar{q}g$, and $gg \rightarrow gg$. We have obtained the relaxation times τ_q and τ_g for the reactions involving quarks (anti-quarks) and gluons respectively by evaluating the rates of the processes. Then the relaxation times are obtained from the well known formula

$$\frac{1}{\tau_m} = \sum_i \frac{1}{\tau_i} \quad (9)$$

where $m = q$ or g and the sum is over all possible processes. As there are a lot of uncertainties in the parameters used, e.g., we have taken the strong coupling constant $\alpha_s = 0.8$, the densities and the temperatures chosen are again arbitrary, the quantitative nature of the curves therefore should not be taken too seriously. However, we expect that the qualitative nature of the curves will not be changed if one uses more exact input parameters.

The figure shows that at low and moderate values of color magnetic field, the shear viscosity coefficient is almost insensitive with the variation of magnetic field strength. However, beyond some upper limit, when the quarks (anti-quarks) occupy only a few low lying Landau levels, the shear viscosity decreases and in the extreme case, when only the zeroth Landau levels are occupied, it exactly vanishes and as a consequence the RHIC sQGP will behave like a super-fluid. We believe that the strength of color magnetic field will be several orders of magnitude larger than the QED quantum critical value for electron and as an outcome, the system should behave like a perfect liquid or super fluid in the extreme case. Then following Feynman we can say that QGP in the extreme case will behave like dry water. We further expect that although the system is a perfect fluid, it can not be a good color conductor. Just like the shear viscosity, the color conductivity will either vanish or will be very small in presence of strong color magnetic field. As a consequence the physical nature of the system will be something like a perfect fluid of bad color conductor or in the extreme case a color neutral super fluid, i.e., the system can not behave like a color super conductor (super fluid of particle with color charges). This will be of course the most surprising phenomenon ever observed. Perhaps such strange physical scenario will be another direct proof of strongly coupled collective behaviour of RHIC sQGP. We further believe that in LHC also the same qualitative nature will be prevailed.

Finally we expect that although RHIC sQGP is a deconfined thermalized strongly correlated quark matter system having collective behaviour, the chiral symmetry will not be restored in the system. The presence of strong color magnetic field will generate mass dynamically just like the QED case.

- [1] J.D. Bjorken, Phys. Rev. **D27**, 140 (1983); S. Chakrabarty and D. Syam, Lett. Nuovo Cim., **41**, 381 (1984); D.K. Srivastava, Jan-e Alam, S. Chakrabarty, B. Sinha and S. Raha, Ann. Phys., **228**, 104 (1993); Y. Akase, M. Mizutani, S. Muroya and M. Yasuda, Prog. Theo. phys., **85**, 305 (1991); A. Dumitru and S. Bass, nucl-th/0007065; T. Hirano and M. Gyulassy, Nucl. Phys. **A769**, 71 (2006); R. Baier, P. Ramatschke and Urs Achim Wiedemann, Phys. Rev. **C73**, 064903 (2006).
- [2] J.H. Applegate, C.J. Hogan and R.J. Scherrer, Phys. Rev. **D35**, 1151 (1987); M.L. Olsen and J. Madsen, Phys. Rev. **D47**, 2313 (1993); J. Kapusta, astro-ph/0101516; D.J. Schwarz, Annalen. Phys. **12**, 220 (2003).




FIG. 1: Variation of shear viscosity coefficient with color magnetic field B normalized with respect to the quantum critical value for electron. Curve (a): $T = 150\text{MeV}$ and $n_B = 5n_0$, Curve (b): $T = 150\text{MeV}$ and $n_B = 8n_0$, and Curve (c): $T = 200\text{MeV}$ and $n_B = 5n_0$,

- [3] Belle Dume, Physics Web, News for April 2005; STAR Collaboration: J. Adams et. al., Nucl. Phys. **A757**, 102 (2005); PHOBOS Collaboration: B.B. Back et. al., Nucl. Phys. **A757**, 28 (2005); PHENIX Collaboration: K. Adcox et. al., Nucl. Phys. **A757**, 184 (2005); BRAHMS Collaboration: I. Arsene et. al., Nucl. Phys. **A757**, 1 (2005).
- [4] M. Riordan and W. A. Zajc, Scientific American, May 2006; D. Teaney, Phys. Rev. **C68**, 034913 (2003); P.K. Kovtun, D.T. Son and A.O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005); G. Policastro, D.T. Son and A.O. Starinets, Phys. Rev. Lett. **87**, 081601 (2001).
- [5] M. Asakawa, S.A. Bass and B. Muller, hep-ph/0603092; see also A. Peshier and W. Cassing, Phys. Rev. Lett. **94**, 172301 (2005).
- [6] S. Chakrabarty, D. Bandopadhyay and S. Pal, Phys. Rev. Lett., **78**, 2898 (1997); D. Bandopadhyay, S. Chakrabarty and S. Pal, Phys. Rev., Lett. **79**, 2176 (1997); Sutapa Ghosh, Soma Mandal and Somenath Chakrabarty, Ann. Phys.**312**, 398 (2004); Soma Mandal, Sutapa Ghosh and Somenath Chakrabarty, Phys. Rev. **C74**, 015801 (2006).
- [7] S.R. de Groot, W.A. van Leeuwen and Ch. G. van Weert, Relativistic Kinetic Theory, North Holland, Amsterdam, 1980.